



Seroprevalence of Newcastle disease virus antibodies in village chickens in the three senatorial zones of Plateau State, Nigeria

AG Balami^{1*}, YS Wungak², SI Bata³ & S Gang³

^{4.} Department of Veterinary Medicine, University of Maiduguri, Nigeria

^{5.} Viral Research Department, National Veterinary Research Institute, Vom, Plateau State, Nigeria

^{6.} Department of Veterinary and Science Laboratory Technology, Federal College of Animal Health and Production Technology, Vom, Plateau State, Nigeria

*Correspondence: Tel.: +2348067703103; E-mail: talktoarrow@yahoo.com

Copyright: © 2022 Balami *et al.* This is an open-access article published under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The most common cause of death in village chickens is Newcastle disease (ND). A study was conducted to evaluate the seroprevalence of ND in the three senatorial zones of Plateau State using the haemagglutination inhibition (HI) test. Blood samples were collected from 389 village chickens of various ages and sexes that were maintained on extensive system of management in nine Local Government Areas (LGAs) across the three Senatorial zones. The overall seroprevalence of ND virus antibodies in village chickens was found to be 36.8% which differed statistically ($p > 0.05$) between LGAs, with the highest prevalence of 8.7% (CI: 6.3–11.9) and lowest prevalence of 1.5% (CI: 0.7–3.3) in Bokkos and Qua'an Pan LGAs, respectively. Also, seroprevalence of NDV antibodies differed significantly ($p > 0.05$) between the three senatorial zones, with 17.5% (CI: 14.0-21.6) in the Northern senatorial zone and 6.7% (CI: 4.6-9.6) in Southern senatorial zone. In the three senatorial zones of Plateau State, this study found prevalence of ND virus antibodies in village chickens. There is a need to increase awareness regarding the importance of vaccinating village chickens to protect them.

Publication History:

Received: 22-12-2021

Revised: 28-04-2022

Accepted: 30-04-2022

Keywords: Newcastle disease, Plateau state, Senatorial zones, Seroprevalence, Village chickens

Introduction

Newcastle disease (ND) is one of the most common poultry diseases worldwide, and it has been identified as one of the leading causes of death among village chicken (Rinle *et al.*, 2019; Egbuji *et al.*, 2017; Lawal *et al.*, 2016). Newcastle disease affects domestic poultry, aviaries, and wild birds and is extremely

contagious. The Newcastle diseases virus strains were divided into three pathotypes by Aldous & Alexander (2010): i.e highly virulent (velogenic), intermediate (mesogenic), and non-virulent (lentogenic), with clinical symptoms ranging from neurologic to pulmonary signs (Alexander *et al.*, 1985). Since its first

report in Ibadan, Nigeria in 1953, ND has grown to become the most important viral disease of chickens, widely recognized by both village chickens and commercial poultry farmers (Lawal *et al.*, 2015), and is widely spread throughout Nigeria, with annual epidemics in highly susceptible poultry flocks (Aliyu *et al.*, 2015).

Village poultry species, especially when grown together or in close vicinity, could play a key role in the epidemiology and spread of the virus to more vulnerable commercial exotic chickens or immune deficient village poultry species (Egbufi *et al.*, 2017). Since Nigeria has a poultry population of 140 million, with backyard poultry production accounting for more than 60% of the total flock and an asset value of more than 5.75 billion dollars (Nnadi & George, 2010), the disease could pose a threat to both successful village poultry and commercial exotic chicken production (Lawal *et al.*, 2016; Egbufi *et al.*, 2017). Although the presence of specific antibodies to ND virus in bird's serum provides little information about the infecting strain of the virus and hence has limited diagnostic value regarding strain identification, the evidence of infection may be sufficient for immediate intervention and control measures (Musa *et al.*, 2009). Several authors studied the seroprevalence of ND in Plateau State (Musa *et al.*, 2009; Egbufi *et al.*, 2017) with no information on the seroprevalence in village chickens in the three Senatorial zones of the state. As a result of these, the aim of this study was to determine the seroprevalence of ND in the three Senatorial zones of Plateau State, Nigeria.

Materials and Methods




Figure 1: Map of Plateau State showing the sampling areas (© Plateau State Government)

Study location

The research was conducted in Plateau State, Nigeria, which is situated between the latitudes of 8° and 10° North and the longitudes of 7° and 11° East. The State has borders with Bauchi State to the east, Taraba State to the South east, Nasarawa State to the South and Kaduna State to the North. It has an average annual rainfall of 1,317 mm in the Southern part and 1460 mm on the high plateau, with an average ambient temperature of 24°C and a relative humidity of 50%. The state's population is estimated to be around 3.5 million people, including more than 30 ethnic groups. Plateau State is divided into three Senatorial zones (NPC 2006; NRSC 2008; NFDP 2009).

Study design

The sampling areas were chosen using a multi-stage sampling technique based on the State's political division into three senatorial zones (Figure 1). Three LGAs were chosen in each senatorial zone using a non-probability sampling technique, and two districts were chosen in each LGA based on the convenience and availability of the live bird market. The LGAs used in this study were Jos North, Jos South, and Jos East in the northern zone, Bokkos, Kanke, and Mangu in the central zone, and Langtang North, Shandam, and Qua'an Pan in the southern zone. In total, 389 blood samples were collected from village chickens across the three Senatorial zones.

Sample size determination

Using the formula $N = Z^2 pq/l^2$, the sample size was calculated (Thrusfield, 2007); where $Z = 1.96$; $P =$

36.4% prevalence in chickens in Zaria (Sule *et al.*, 2019); $q = 1 - p$; $l = 5.0\%$ allowable error. There were 198 samples in all. But, 389 village chickens from the nearby area were sampled,

Blood sample collection

Blood sample was collected from each of the village chickens, irrespective of age and sex with history of no vaccination against ND. A 21 G needle attached on a 5ml syringe was used to collect two millilitres of blood from the wing vein and allowed to coagulate in the syringe. For complete serum separation, the clotted blood was left at room temperature overnight. The separated serum was then decanted into cryovials, immediately delivered to the laboratory and stored at -20°C until it was analysed for ND antibodies.

Serology

The harvested serum samples were tested using haemagglutination and haemagglutination inhibition tests. According to the method of the Office International des Epizootics (OIE, 2010), 1% suspension of chicken red blood cells (RBCs) was prepared as an indicator for the haemagglutination (HA) and haemagglutination inhibition (HI) tests (OIE, 2010). According to the method of Allan and Gough (1974), HA titre of a standard NDV antigen was determined, which was diluted to contain 4HA units. In an HI test, ND antibody titres were determined in harvested sera using reconstituted antigen containing the 4HA units. The agglutination was assessed by tilting the plates. Only those wells in which the RBCs streamed at the same rate as the control wells were considered to show HI.

Data analysis

Data were stored in Microsoft excel spread sheet. Descriptive statistics was carried out using Microsoft excel and proportions were obtained using open Epi version 2.3.1 statistical tool (Open-source epidemiological statistic for public health calculation). Chi square was used to measure the

association between location and sex and the prevalence of ND antibodies. Value of $p < 0.05$ were considered significant. Results were presented in tables and graphs.

Results

The result of this study revealed an overall Seroprevalence of 36.8% (CI: 32.1-41.6) in the three Zones under the study. The Seroprevalence of ND virus antibodies in the village chickens in the selected LGA's in the three Senatorial zones showed the highest prevalence of 8.7% (CI: 6.3 – 11.9) in Bokokos, (Central Zone). This was followed by 7.9% (CI: 5.7 – 11.1) in Jos-South (Northern Zone) and 1.5% (CI: 0.7– 3.3) in Qua'an Pan (Southern zone) with statistically significant differences $P < 0.05$ (Table 1).

Statistically significant difference ($p < 0.05$) was observed in the Seroprevalence of NDV antibodies between the three Senatorial zones. The Northern Senatorial zone had the highest Seroprevalence of 17.5% (CI: 14.0 - 21.6), followed by the Central zone; 12.6% (CI: 9.7 - 16.3) and the Southern zone with 6.7% (CI: 4.6 - 9.6) (Table 2).

There was no statistically significant difference ($p > 0.05$) in the Seroprevalence of ND virus

Table 1: Seroprevalence of Newcastle disease virus antibodies in three Senatorial zones of Plateau State based on Local Government Areas

LGA	Number Examined	Number Positive	Number Negative	Proportion %	Confidence Interval
Northern Senatorial Zone					
Jos South	48	31	17	7.9	5.7 - 11.1
Jos East	44	24	20	6.2	4.2 - 9.0
Jos North	44	13	31	3.34	1.9 - 5.6
Central Senatorial Zone					
Bokokos	48	34	14	8.7	6.3 - 11.9
Kanke	24	5	19	1.3	0.6 - 2.9
Mangu	43	10	33	2.6	1.4 - 4.7
Southern Senatorial Zone					
Langtang N	47	13	34	17.5	14.0 - 21.6
Shendam	45	7	38	12.6	9.7 - 16.3
Qua'an Pan	46	6	40	6.7	4.6 - 9.6

Chi square =74.43, P value = < 0.0000001 ($P < 0.05$), Degree of freedom =8
 Overall prevalence =36.8% (CI: 32.1-41.6)

Table 2: Seroprevalence of Newcastle disease virus antibodies based on three Senatorial zones of Plateau State

Zone (Z)	Number Examined	Number Positive	Number Negative	Proportion %	Confidence Interval
Northern Z.	136	68	68	17.5	14.0 - 21.6
Central Z	115	49	66	12.6	9.7 - 16.3
Southern Z.	138	26	112	6.7	4.6 - 9.6
Total	389	143	246		

Chi square =31.01, P value = 0.00000185 ($P < 0.05$), Degree of freedom =2

Table 3: Seroprevalence of Newcastle disease virus antibodies in three Senatorial zones of Plateau State based on sex

Sex	Number Examined	Number Positive	Number Negative	Proportion %	Confidence Interval
Males	194	71	123	18.3	14.7 - 22.4
Females	195	72	123	18.5	14.9 - 22.7
Total	389	143	246		

Chi square = 0.0044, P value = 0.95, (P < 0.05), Degree of freedom = 1

antibodies between the sexes of the village chickens. A Seroprevalence of 18.5% (CI: 14.7 – 22.4) was obtained for females while the value for males was 18.3% (CI: 14.7 – 22.4) (Table 3).

Discussion

The overall Seroprevalence rate of 36.8% recorded in this study poses a significant danger to effective village chicken production (Lawal *et al.*, 2016) and even more so to Plateau state's commercial poultry industry. If not appropriately controlled, the 36.8% Sero-positive chickens could serve as a source of infection for a vast population of village and commercial poultry. This is because most infected birds might shed substantial amounts of ND virus in their faeces (Alexander *et al.*, 1985), which may be spread by people, equipment, vehicles, contaminated poultry and poultry products, contaminated feed and water, and other animals (Capua & Alexander, 2009). The overall seroprevalence recorded in this study is in contrast to the report of Musa *et al.* (2009) who determined a seroprevalence of 51% in Plateau State. The reduction may be attributed to the constant awareness created by veterinarians on the need to vaccinate village chickens with the ND I₂ vaccine, which is readily available at an affordable price. The ND antibodies detected in the village chickens might have been as a result of natural exposure to the ND virus, since the chickens had no history of previous vaccination against ND.

Although, Musa *et al.* (2009) based their data on only four LGAs (2 each from the North and Southern zones of Plateau State), the total seroprevalence rate of NDV in village hens observed in this study does not correspond with the 51% seroprevalence rate reported by Musa *et al.* (2009). This investigation also differs from that of Lawal *et al.* (2016), who found 62.7% seroprevalence to ND virus antibodies in village hens in Gombe State, and Nwanta *et al.* (2008), who found 32.2% in Kaduna State. Other studies, such as those by Ameji *et al.* (2011) and Chollum *et al.* (2013), Jibril *et al.* (2014), and Eze & Ike (2015), revealed seroprevalence rates of 96%, 35.8%, 25.5% and 65.1% respectively.

The differences between the results of this study and those of Ameji *et al.* (2011), Chollom *et al.* (2013), Jibril *et al.* (2014), and Eze & Ike (2015) was that the latter collected samples from village chickens in live bird markets, where birds were known to be brought from various locations, including neighbouring states and congregate for sales.

The significant difference between the LGAs could be due to ecological differences in the ND virus's activity, the effect of the environment on the virus's viability and spread (Orajaka *et al.*, 1999; Eze & Ike 2015), or the high concentration of commercial poultry farms in the LGAs sampled in Plateau state's Northern and Central zones. The large volume of manure produced in concentrated poultry production areas can contribute to the contamination of ground and surface waters (Mallin & Cahoon, 2003).

Manure from poultry litter is generally used as fertilizer and applied to fields for growing agricultural commodities (USPEA, 1998). Jos South and Bokkos LGAs, which are in the State's northern and central zones, are known to have a large concentration of maize and Irish potato farmers. These agricultural products are grown using organic manure from poultry litters, which could be another reason for the significantly higher seroprevalence rate observed among the village chickens sampled in the Northern and Central zones (where high concentration of commercial poultry farms are located), when compared with the Southern zone of the State. This is because ND viruses are frequently discharged in high amounts in poultry faeces (Alexander *et al.*, 1985). Village chickens are known to scavenge the environment for food, and ingestion of faeces during such activity could be a major route of transmission of ND virus among them (Sa'idu *et al.*, 2006).

The insignificant difference observed in the seroprevalence of ND virus between sexes showed that ND virus has no sex specificity and therefore can infect both sexes of village chickens and can serve as reservoir to ND virus. This result agrees with that of Elifuraha & Emmanuel (2021), who also reported no significant difference in the seroprevalence of ND virus between both sexes of village chickens in

Njombe and Bahi Districts in Tanzania. But however, they differ from the reports of Unigwe *et al.* (2020), who reported a higher seroprevalence of ND virus in males than in females in Ido and Atiba LGA's of Oyo State, Nigeria, and Conteh *et al.* (2020), who showed a significantly higher seroprevalence of ND virus in females than in males in Sierra Leone.

In conclusion, the study found a prevalence of ND virus antibodies in village chickens. As a result, it is recommended that more awareness be raised about the importance of vaccinating village chickens to protect them against ND virus.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

Aldous EW & Alexander DJ (2010). Technical review: Detection and differentiation of Newcastle disease virus (avian paramyxovirus type 1). *Avian Pathology*, doi.10.1080/03079450120044515.

Alexander DJ, Wilson GWC, Russell PH, Lister SA & Parsons G (1985). Newcastle disease outbreaks in fowl in Great Britain during 1984, *Veterinary Records*, **117**(17): 429-434.

Aliyu HB, Saidu L, Abdu PA & Oladele SB (2015). A retrospective analysis of Newcastle disease diagnosed at poultry clinic of Ahmadu Bello University, Zaria, Nigeria. *Sokoto Journal of Veterinary Science*, **13**(3):42-48.

Allan WH & Gough E (1974). A standard haemagglutination inhibition test for Newcastle disease a comparism of macro and micro methods. *Veterinary Record*, doi.10.1136/vr.95.6.120.

Ameji ON, Abdu PA & Saidu L (2011). Seroprevalence of avian influenza and Newcastle disease and Gumboro disease in chicken in Kogi State, Nigeria. *Bulletin of Animal Health and Production in Africa*, **59**(4): 411-418.

Capua I & Alexander DJ (2009). Avian Influenza and Newcastle Disease A Field and Laboratory Manual. Milan: Springer-Verlag. Pp 19-21.

Chollom SC, Emerhirhi FT, Akwaowo EE, Ogbaji JU & Fyaktu EJ (2013). Implication of Newcastle disease virus in local chickens at live bird markets in Jos, Nigeria. *International Journal of Current Research and Review*, **5**(14): 2872-2874.

Conteh AM, Moiforay SK, Sesay ME & Kallon S (2020). Prevalence of Newcastle disease virus

antibodies in apparently healthy chickens in Sierra Leone. *Animal and Veterinary Sciences*, doi.10.11648/j.av.s.20200805.12.

Egbuji AN, John OI, Shittu IA, Echeonwu GON, Uwanibe EL, Bwala D, Kiagama J, Dung E, Nyam D, Obene C, Kwanga A, Mukaila F, Dagwong H, Lokason S, Useh N & Rabo J. (2017). Screening of rural scavenging birds for the presence of detectible protective Newcastle disease antibodies in some selected rural communities of Plateau State. *African Journal of Microbiology Research*, **11**(37): 1431-1433.

Elifuraha BM & Emmanuel B (2021). Seroprevalence of Newcastle Disease Virus in Local Chickens in Njombe and Bahi Districts in Tanzania. *Research Square*, doi: doi.10.21203/rs.3.rs-712858/v1

Eze IA & Ike AC (2015). The serological status for Newcastle disease in local chickens of live bird markets and households in Nsukka, Enugu State, Nigeria. *Nigeria Journal of Microbiology*, **29**(2): 3096-3104.

Jibril AH, Umoh JU & Kabir J (2014). Newcastle disease in local chickens of live bird markets and households in Zamfara State, Nigeria. *International Scholarly Research Notices*, doi.10.1155/2014/513961.

Lawal JR, El-Yuguda AD & Ibrahim UI (2016). Survey on Prevalence of Newcastle Disease Antibodies in Village Poultry at Live Birds Markets in Gombe, Nigeria. *Journal of Animal Science and Livestock Production*, doi.10.21767/2577-0594.100001.

Lawal JR, Jajere SM, Mustapha M, Bello AM, Wakil Y, Geidam YA, Ibrahim UI & Gulani IA (2015). Prevalence of Newcastle Disease in Gombe, Northeastern Nigeria: A Ten-Year Retrospective Study (2004 –2013). *British Microbiology Research Journal*, **6**(6): 367-375.

Mallin MA & Cahoon LB (2003). Industrialized animal production—a major source of nutrient and microbial pollution to aquatic ecosystems. *Population and Environment*, **24**(5): 369–385.

Musa U, Abdu PA, Dafwang II, Umoh JU, Saidu L, Mera UM & Edache JA (2009). Seroprevalence, seasonal occurrence and clinical manifestation of Newcastle disease in rural household chickens in plateau state, Nigeria, *International Journal of Poultry Science*, **8**(2): 200– 204.

- NFDP (2009). National Fadama Development Project: Plateau Agricultural Development Project, Quarterly Review Bulletin. Pp 4-9.
- Nnadi PA & George SO (2010). A cross-sectional survey on parasites of chickens in selected villages in the sub-humid zones of South Eastern Nigeria. *Journal of Parasitology Research*, **4**(1):1-6.
- NPC (2006). National Population Commission. Plateau State Statistics Tables, Head office Abuja, Nigeria. 12-18. Pp 12-18.
- NRSC (2008). National Remote Sensing Corporation Jos. Classified geo sensing data for Plateau State, Nigeria. National Remote Sensing Head Office, Jos, Plateau, Nigeria. Pp 2-7.
- Nwanta JA, Abdu PA & Ezema WS (2008). Epidemiology, challenges and prospects for control of Newcastle disease in village poultry in Nigeria. *World's Poultry Science Journal*, **64**: 119-127.
- OIE (2010). Newcastle Disease, Manual of standard for diagnostic tests and vaccines. Office International des Epizootics, (OIE), Paris. Pp 221-232.
- Orajaka LJE, Adene DF, Anene B & Onuoha EA (1999). Seroprevalence of Newcastle disease in local chickens from Southeast derived Savannah zone of Nigeria. *Rvued'Elevage et de Medicine Veterinaires des Pays Tropicaux*, **52**(3-4): 185-188.
- Rinle KP, Oladele SB, Esievo KAN, Ibrahim NDG, Ahmed JS, Moses GD, Rimfa AG, Choji TPP, Ugbe AD, Anueyiagu DM, Atam SK, Shekaro A & Barde IJ (2019). Clinico-pathological and Immunohistochemistry of Newcastle Disease in Indigenous Chickens and Some Wild Birds in Plateau State, Nigeria. *Journal of Advances in Microbiology*, **16**(4): 1-15.
- Sa'idu L, Abdu PA, Tekdek LB, Umoh JU, Usman M & Oladele SB (2006). Newcastle disease in Nigeria. *Nigerian Veterinary Journal*, **27**(2): 23-32.
- Sule AG, Abdu PA, Junaidu K & Kia GSN (2019). Seroprevalence and detection of Newcastle disease virus matrix gene in domestic local breed of chickens from eight communities in Bauchi State, Nigeria. *Nigerian Veterinary Journal*, **40**(2): 118 – 126.
- Thrusfield M (2007). *Veterinary Epidemiology*, Third edition. Blackwell Science Limited, London. Pp 227–247.
- Unigwe CR, Shobowale OM, Enibe F, Ajayi JO & Koleosho SA (2020). Sero-prevalence of Newcastle disease in apparently healthy normal feathered local chickens in Ido and Atiba local government areas, Oyo State, Nigeria. *Journal of Tropical Agriculture, Food, Environment and Extension*, **19**(4): 37 – 42.
- USPEA (1998). United State Poultry and Egg Association: Environmental framework and implementation strategy for poultry operations. <http://www.poultryegg.org/Environment/STRATEGY1.html>, retrieved 12-09-2021.