

CONTRIBUTIONS OF ENVIRONMENTAL FACTORS TO THE ESTIMATIONS OF *EUSTRONGYLIDES AFRICANUS* LARVAE DENSITIES IN *CLARIAS GARIEPINUS* AND *CLARIAS ANGUILLARIS* FROM BIDA FLOOD PLAIN OF NIGERIA

T.I.I. IBIWOYE¹*, R.A. OGUNSUSI², A.M. BALOGUN³ AND J.J. AGBONTALE¹

¹Aquatic Pathobiology Programme, National Institute for Freshwater Fisheries Research
P.M.B. 6006, New Bussa, Niger State, Nigeria

²Department of Animal Production and Health, Federal University of Technology, P.M.B. 704, Akure, Ondo State, Nigeria

³Department of Fisheries and Wildlife, Federal University of Technology, P.M.B. 704, Akure, Ondo State, Nigeria

*Correspondence Author

Abstract

The contribution of environmental factors to the estimations of *Eustrangylides africanus* larvae densities in *Clarias gariepinus* and *C. anguillaris* from Bida flood plain of Nigeria was investigated. Six environmental factors (sand, silt-clay, soil pH, water turbidity, dissolved oxygen and total phosphate content) having positive or negative correlation coefficient (*r*) between 0.50 and 0.85 contributed to the estimations of *E. africanus* larvae densities in the wild population of *Clarias* species.

Introduction

Disease aetiology is a triad complex (Snieszko, 1974), which includes the host (fish), the parasite (agent), and the micro – and macro – habitats (environment). Climate is the average of atmosphere conditions (that is desirable minimum of 35 years) involving the systematic observation, recording and processing of the various elements: temperature, humidity, sunshine, evaporation, rainfall and others (Areola *et al.* 1992). Physiographically, the study area consists of level of gentle rolling plains. The monotonous of the landscape is only broken by some conical or flat-topped residual hills. The elevation of the plain is between 120 and 240m above sea level (Adeleye, 1976). Meteorologically, the area has a marked wet (April to October) and dry (November to May) season and lies within the 1270mm isohyets (Kowal and Knabe, 1972). The soil is reddish and highly leached with low base status, consisting of loamy sand over sand-clay to clay subsurface horizons (Esu, 1976) of the Nupe sandstone with ustic moisture response and in isohyperthermic temperature regime.

Several investigations indicated that the distribution of ecto-and endo-parasites in the fish host populations are related to their environment and their geographical locations (Toweill and Tabor 1982; Bates and Kennedy, 1991). The occurrences of helminth parasites in Nigerian fishes have been extensively studied (Alfred-Okya, 1989; Emere, 2000) were not related to contribution of environmental factors. Thus changes brought about by the environmental, fish and invertebrate host populations, and helminth parasites incidences might not be understood. *Clarias* are highly prized and are found all year round in markets of Bida and its environs. The aim of this study therefore is to investigate the contribution of some environmental factors to the estimation of *Eustrangylides africanus* larvae in *Clarias* species from Bida floodplain of Nigeria. This study will provide base line information on the contributions of environmental factors to the estimation of *Eustrangylides africanus* larvae densities in *Clarias* species from Bida floodplain of Nigeria.

Materials and Methods

The study covered an area between longitude 5° 45' to 6° 15'E and latitude 8° 30' to 9° 10'N within the southern Guinea savannah zone of Nigeria (Areola *et al.* 1991). Regular field data were collected from the fishing localities (Doko, Dokogi, Fokpo and Dutsu) in Bida floodplain on the soil, water and fish components.

The rapid partial analysis of soil involved splitting of the samples into a sand fraction (particles greater than 62µ) and a silt-clay fraction (particles less than 62µ) according to Wentworth scale as modified by Krumben (Buchanan and Kain, 1971). Soil hydrogen concentration pH in water (1:1 soil to water) was measured by inserting the pH meter (Model 264A, Jenway, England) electrodes into partly settled suspension stirred during measurement.

Water samples were collected, with a 2-litre van-Dorn water sampler, into clean plastic container and preserved for respective laboratory analysis carried out. The water temperature was measured on the spot by using graduated mercury in glass thermometer calibrated at 0.2°C interval with a range of 0 to 40°C, recorded immediately for each water sample into sampled bottles. The conductivity, turbidity and total dissolved solids were measured in their respective units by standard methods described by HACH (1980). The pH of the water samples was determined by the colorimeter method described HACH (1980) using the Lovibond pH comparator with bromothymol blue as indicator. The acid modification of Winkler's iodometric method described by ALPHA (1989) involving the use of four major reagents (manganous sulphate Mn+2SO₄): sodium of potassium hydroxide and iodide (Winkler's reagents); concentrated sulphuric acid (H₂SO₄) and sodium thiosulphate (Na₂SO₂O₃) estimate the dissolved oxygen content of the water sample in mg/l. The nitrate and total phosphate content was measured in mg/l in each water sample using the cadmium reduction and ascorbic acid method respectively as described by HAC (1980).

The records of the climatic factors (minimum and maximum temperature, relative humidity, rainfall, evaporation and sunshine) kept between January 1959 and December 1999 at the Metrological Unit of the National

Cereal Research Institute (NCRI) Badeggi – Bida were assembled and analyzed as secondary data.

Routine examination were carried out on four hundred and eighty specimens of *Clarias gariepinus* and *C. anguillaris* of different sexes, lengths and weights. The specimens were randomly sampled from four fishing localities of Bid floodplain species to determine occurrence of *Eustrongylides africanus* larvae in relation to sex and season of the year as described by Margolis *et al.* (1982). Simple linear regression/correlation analysis were carried out to examine any associations among prevalence, mean intensity and abundance of *Eustrongylides africanus* larvae in *Clarias* species of Bida floodplain and the sixteen environmental factors [maximum temperature (x1), minimum temperature (x2) and relative humidity (x3); rainfall (x4); sunshine (x5), evaporation (x6), sand (x7), silt-

clay (x8), soil pH (x9), water temperature (x10), conductivity (x11), turbidity (x12), total dissolved solids (x13), water pH (x14), dissolved oxygen (x15) and total phosphate content (x16).]

Results and Discussion

The correlated coefficient R for sixteen environmental factors with the occurrences of *Eustrongylides africanus* larvae in *Clarias* species frp, Bida floodplain is shown on Table 1. Six, five and six out of the 52 combinations of the sixteen environmental factors with known correlation coefficient ($r \geq \pm 0.50$) contributed respectively to the estimation of prevalence, mean intensity and abundance of *E. africanus* larvae densities in *Clarias* species (Tables 2, 3 and 4) were sand, sand, silt – clay, soil pH, turbidity, dissolved oxygen and phosphate as follows:

Equations for prevalence

$Y = -145x + 140.7$ ($r = -0.83$) with sand
 $Y = -1.24x + 18.1$ ($r = -0.54$) with silt clay
 $Y = -5.9x + 65$ ($r = -0.50$) with soil pH
 $Y = -0.55x + 24.5$ ($r = -0.51$) with turbidity
 $Y = -0.52x + 48.2$ ($r = -0.69$) with dissolved oxygen
 $Y = -0.03x + 35.1$ ($r = -0.70$) with total phosphate content

Equations for mean abundance

$Y = -0.05x + 4.8$ ($r = -0.85$) with sand
 $Y = -0.06x + 0.2$ ($r = -0.76$) with silt clay
 $Y = -0.28x + 2.4$ ($r = -0.65$) with soil pH
 $Y = -0.02x + 1.3$ ($r = -0.59$) with dissolved oxygen
 $Y = -0.04x + 1.0$ ($r = -0.79$) with total phosphate content

Equations for mean intensity

$Y = -0.8x + 8.4$ ($r = -0.75$) with sand
 $Y = -0.12x + 1.2$ ($r = -0.85$) with silt clay
 $Y = -0.45x + 5.0$ ($r = -0.63$) with soil pH
 $Y = -0.07x + 2.7$ ($r = -0.80$) with total phosphate content

The silt-clay of the soil and water turbidity were positively correlated to the prevalence of *E. africanus* larvae in *Clarias* species. Thus silt-clay of the soil and water turbidity increase as the prevalence of *E. africanus* larvae species increases and vice versa. The sand, soil pH dissolved oxygen and total phosphate content were negatively correlated to the prevalence of *E. africanus* larvae in *Clarias* species. Thus the sand, soil pH dissolved oxygen and total phosphate content would increase as the prevalence of *E. africanus* larvae species in *Clarias* species increase, and vice versa.

The silt-clay of the soil was positively correlated to the mean intensities of *E. africanus* larvae in *Clarias* species. Thus silt-clay of the soil increases as the main intensity of *E. africanus* larvae species increases and vice versa. The sand, soil pH and total phosphate content were negatively

correlated to the mean intensities *E. africanus* larvae in *Clarias* species. Thus the sand, soil pH and total phosphate content increases as the mean content increases as the mean intensities of *E. africanus* larvae species in *Clarias* species decrease, and vice versa.

The silt-clay of the soil was positively correlated to the mean abundance of *E. africanus* larvae in *Clarias* species. Thus, silt-clay of the soil increases as the main abundance of *E. africanus* larvae species increases, and vice versa. The sand, soil pH, dissolved oxygen and total phosphate content were negatively correlated to the mean abundance of *E. africanus* larvae in *Clarias* species. Thus, the sand, soil pH, dissolved oxygen and total phosphate content increases as the mean content increases as the mean abundance of *E. africanus* larvae species in *Clarias* species decrease, and vice versa.

Table 1: Correlation coefficient (*r*) for sixteen environmental factors with occurrence of *Eustrongylides africanus* larvae infection in *Clarias* species from Bida Floodplain

	Preva.	Inten	Abun.	Max Tem (x ₁)	Min Tem (x ₂)	RH (x ₃)	Rain (x ₄)	Evapo (x ₅)	Sun (x ₆)	Sand (x ₇)
Prevalence	1									
Intensity	0.82	1								
Abundance	0.92	0.97	1							
Temp max (x ₁)	0.14	0.07	0.13	1						
Temp min (x ₂)	-0.19	-0.09	0.18	-0.23	1					
RH (x ₃)	-0.38	-0.27	-0.38	-0.85	0.65	1				
Rain (x ₄)	-0.03	0.01	-0.06	-0.85	0.67	0.91	1			
Evapo (x ₅)	0.09	0.03	0.03	0.40	0.76	0.01	0.14	1		
Sun (x ₆)	-0.04	-0.20	-0.14	0.64	-0.50	-0.61	-0.82	-0.15	1	
Sand (x ₇)	-0.83	0.75	0.85	0.45	-0.35	-0.69	0.36	-0.12	-0.05	1
Silt-clay (x ₈)	0.83	0.75	0.85	0.45	-0.35	-0.69	0.36	-0.12	-0.05	-1.00
pH (x ₉)	-0.49	-0.63	-0.65	-0.66	0.33	0.79	0.58	-0.17	-0.14	0.87
Temp (x ₁₀)	0.16	0.20	0.15	-0.05	0.66	0.28	0.41	0.64	-0.28	0.21
Cond. (x ₁₁)	0.29	0.09	0.11	-0.25	-0.03	0.22	0.13	-0.22	0.39	0.16
Turb (x ₁₂)	0.51	0.31	0.33	-0.40	0.47	0.44	0.61	0.29	-0.22	-0.10
TDS (x ₁₃)	0.41	-0.09	0.06	-0.33	0.04	0.22	0.35	-0.02	0.02	0.02
pH (x ₁₄)	0.36	0.01	0.16	0.81	-0.41	-0.81	-0.76	0.20	0.78	-0.47
DO (x ₁₅)	-0.069	-0.45	-0.59	-0.71	0.54	0.88	0.70	-0.04	-0.54	0.74
PO ₃ (x ₁₆)	-0.70	-0.80	-0.79	-0.46	0.25	0.54	0.43	-0.04	-0.38	0.62

X1 – Maximum temperature, X2 – Minimum temperature, X3 – Relative humidity, X4 – Rainfall, X5 – Evaporation, X6 – Sunshine, X7 – Sand, X8 – Silt-Clay, X9 – Soil pH, X10 – Water Temperature, X11 – Conductivity, X12 – Turbidity, X13 – Total dissolved solids, X14 – Water pH, X15 – Dissolved oxygen, X16 – Total phosphate content.

Table 1 (continued): Correlation coefficient @ for sixteen environmental factors with occurrence of *Eustrongylides africanus* larvae infection in *Clarias* species from Bida Floodplain

	Silt/ Clay (x ₈)	pH (x ₉)	Tem (x ₁₀)	Con. (x ₁₂)	Tur (x ₁₂)	TDS (x ₁₃)	pH (x ₁₄)	DO (x ₁₅)	PO ₃ (x ₁₆)
Silt-clay (x ₈)	1								
pH (x ₉)	-0.87	1							
Temp (x ₁₀)	-0.21	-0.12	1						
Cond. (x ₁₁)	-0.16	0.47	0.01	1					
Turb (x ₁₂)	0.10	0.30	0.61	0.67	1				
TDS (x ₁₃)	-0.02	0.50	0.05	0.67	0.72	1			
pH (x ₁₄)	0.47	-0.41	-0.09	0.18	-0.05	0.23	1		
DO (x ₁₅)	-0.74	0.64	0.26	-0.11	0.08	-0.15	-0.84	1	
PO ₃ (x ₁₆)	-0.62	0.60	0.03	-0.33	-0.15	0.09	-0.42	0.71	1

X1 – Maximum temperature, X2 – Minimum temperature, X3 – Relative humidity, X4 – Rainfall, X5 – Evaporation, X6 – Sunshine, X7 – Sand, X8 – Silt-Clay, X9 – Soil pH, X10 – Water Temperature, X11 – Conductivity, X12 – Turbidity, X13 – Total dissolved solids, X14 – Water pH, X15 – Dissolved oxygen, X16 – Total phosphate content.

Table 2: Estimations of prevalence of *Eustrongylides africanus* larvae infection in *Clarias* species from Bida Floodplain using environmental factors with known correlation coefficients ($r \geq \pm 0.50$)

Environmental Factors (EFs)	Equation for Prevalence (Y)	Equivalence of EFs (When Y = 0)	Equivalence of (EFs) (For 1 % Change in Y)
Sand (%)	$Y = -145 x_1 + 140.7$ ($r = -0.83$)	$x = -96.7$	$x = -0.7$
Silt Clay (%)	$Y = -1.24 x_2 + 18.1$ ($r = -0.54$)	$x = -14.6$	$x = 0.8$
Soil pH (unit)	$Y = -5.9 x_3 + 65$ ($r = -0.50$)	$x = 110.1$	$x = -1.7$
Turbidity (mg/l)	$Y = -0.55 x_4 + 24.5$ ($r = -0.51$)	$x = -444.7$	$x = 8.2$
Dissolved Oxygen (mg/l)	$Y = -0.52 x + 48.2$ ($r = -0.69$)	$x = 92.6$	$x = 0.9$
Phosphate (mg/l)	$Y = -0.03 x_5 + 35.1$ ($r = -0.70$)	$x = 1252.5$	$x = 5.7$

Table 3: Estimations of mean intensity of *Eustrongylides africanus* larvae infection in *Clarias* species from Bida Floodplain using environmental factors with known correlation coefficients ($r \geq \pm 0.50$)

Environmental Factors (EFs)	Equation for Intensity (Y = No of larvae per infected fish sampled)	Equivalence of EFs (When Y = 0)	Equivalence of (EFs) (For 1 % Change in Y)
Sand (%)	$Y = -0.079 x_1 + 8.39$ ($r = -0.75$)	$x = 106.2$	$x = -12.8$
Silt Clay (%)	$Y = -1.2 x_2 + 1.23$ ($r = -0.85$)	$x = -10.3$	$x = 8.4$
Soil pH (unit)	$Y = -0.45 x_3 + 5.04$ ($r = -0.63$)	$x = 11.2$	$x = -2.2$
Phosphate (mg/l)	$Y = -0.55x + 24.5$ ($r = -0.51$)	$x = 38.0$	$x = -6.0$

Table 4: Estimation of mean abundance of *Eustrongylides africanus* larvae infection in *Clarias* species from Bida Floodplain using environmental factors with known correlation coefficients ($r \geq \pm 0.50$)

Environmental Factors (EFs)	Equation for Prevalence (Y)	Equivalence of EFs (When Y = 0)	Equivalence of (EFs) (For 1 % Change in Y)
Sand (%)	$Y = -0.053 x_1 + 4.80$ ($r = -0.85$)	$x = 90.6$	$x = -18.9$
Silt Clay (%)	$Y = 0.062 x_2 + 0.17$ ($r = 0.76$)	$x = -2.7$	$x = 10.7$
Soil pH (unit)	$Y = -0.28 x_3 + 2.42$ ($r = -0.65$)	$x = 8.6$	$x = -3.7$
Dissolved Oxygen (mg/l)	$Y = -0.016 x_4 + 1.33$ ($r = -0.59$)	$x = -83.1$	$x = -62.5$
Phosphate (mg/l)	$Y = -0.016 x_4 + 1.33$ ($r = -0.59$)	$x = 23.1$	$x = -23.8$

Conclusion

Many fish farmers, as well as hatchery operators, often need to know the health statues so as to make management decisions like how weight gain feed intake and utilization, and also to administer the right damage of medication. Reasonable skill in estimating parasitic infection density is necessary for fishery workers as it will frequently be necessary to estimate parasite density when facilities to measure environmental factors are not readily available or their use are not practicable. This study has shown that environmental factors, as media for proper fish productivity, could be used to estimate the density of parasitic infections in freshwater fishes. Thus, a simple device for quick and accurate estimation of the density of parasites in fishes could be derived for research and development purposes.

Acknowledgements

The authors are grateful to the Federal Government of Nigeria through the Federal Ministry of Agriculture and Rural Development and National Institute for Freshwater Fisheries Research (NIFFR) New Bussa for providing the fund for this study.

References

- Adeleye, D. R. (1976). The geology of the Middle Niger Basin. In: C. A. Kogbe (ed.), *Geology of Nigeria*. Pp. 283-287.
- Alfred-Ockiya, J. F. (1989). Preliminary survey of parasites of *Clarias* in the Niger Delta Area of Nigeria. *Proceedings of the 4th Annual Conference of the Fisheries Society of Nigeria (FISON, Port Harcourt, 26th – 28th Nov. 1985*. pp. 210-215.
- Anosike, J.C.; E. Omoregie; P.C.. Ofojekwu and I. E. Nweke (1992). A survey of helminth parasites of *Clarias gariepinus* in Plateau State, Nigeria. *Journal of Aquatic Science*. 7:39-43.
- APHA (American Public Health Association) (1989). *Standard methods for the examination of water and waste-water (17th edition)*. American Public Health Association, American Water Works Association and Water Pollution Control Federation, Washington, DC. 1391pp.
- Aresola, O; O. Irukebhe; K. Ahmed; B. Adeleke and G. C. Leong (1992). *Certificate Physical and Human Geography for Secondary Schools*. University Press, Plc. Ibadan, 406pp.
- Auta, J; S.J. Oniye and J. A. Adakole (1999). The helminth parasites of the gastro-intestinal tract of Synodontis species in Zaria, Nigeria. *Zuma Journal Pure and Applied Sciences*. 2(2): 47-53.
- Bates, R.M. and C. R. Kennedy (1991). Site availability and density-dependent constraints on the acanthocephalan *Pomphorhynchus laevis* in rainbow trout *Onchorynchus mykiss* (Walbaum), *Parasitology*, 102:405-410.
- Buckannan, J. B. and J. M. Kain (1971). Measurement of the physical and chemical environment. In: *Methods for the study of marine benthos*. Holmes, N. A. & A.D. Melntyre (eds.) IBP Handbook No. 16. Blackwell Scientific Publishers. Pp. 35-36.
- Emere, M. C. (2000). Parasitic infection of the Nile perch *Lates niloticus* (L) in River Kaduna. *Journal of Aquatic Sciences*. 15:51-54.
- Esu, I. E. (1986). Morphology and classification of the soils of Nupe sandstone formation in Niger State, Nigeria. *Samaru Journal of Agricultural Research*. 41(1&2); 13-23.
- Ezenwaji, H.M.G. and P.C.O. Ilozumba (1992). Helminth fauna of four West African small *Clarias* species (Osteichthyes: Clariidae) from Nigeria. *Journal of African Zoology*. 106:391-400.
- Hach (1980). Hach water analysis handbook. Hach Chemical Company, Loveland, Colorado, U. S.A. 302pp.
- Ibiwoye, T.I.L; A.M. Sule; P. U.A. Okojie and J. J. Agbontale (1996). Prevalence of helminth infestation in freshwater fadama fishes of the Bida Area, Nigeria. National Institute for Freshwater Fisheries Research (NIFFR) Annual Report ISSN.0331 – 9296.p.43-49.
- Kowal, J.M. and D. T. Knabe (1972). An agroclimatological atlas of the Northern States of Nigeria with explanatory notes. A.B.U. Press, Zaria, Nigeria. 111pp.
- Margolis, L.; G. W. Esch; J. C. Homes; A. M. Kuris and G. A. Schad (1982). The use of ecological terms in

- parasitology (Report of an ad-hoc committee of the American Society of Parasitologists. *Journal of Parasitology*. **68**:131-133.
- Snieszko, S. F. (1974). The effect of environmental stress on outbreak of infectious diseases of fishes. *Journal of Fish Biology*. **6**:197-208.
- Toweill, D. E. and J. E. Tabor (1982). River otters, *Lutra Canadensis*. In: *Wild mammals of North America, biology, management and economics*. J. A. Chapman and G. A. Feldhammer (eds.) Johns Hopkins University Press, Baltimore, Maryland, p. 688-703.